

## NATIONAL INSTITUTES OF HEALTH — FAT SCREENER

1. Think about your eating habits over the past 12 months. About how often did you eat or drink each of the following foods? Remember breakfast, lunch, dinner, snacks, and eating out. Blacken in only one bubble for each food.

TYPE OF FOOD	Never	Less than Once Per Month	1-3 Times Per Month	1-2 Times Per Week	3-4 Times Per Week	5-6 Times Per Week	1 Time Per Day	2 or More Times Per Day
Cold cereal	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Skim milk, on cereal or to drink	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Eggs, fried or scrambled in margarine, butter, or oil	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Sausage or bacon, regular-fat	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Margarine or butter on bread, rolls, pancakes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Orange juice or grapefruit juice	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Fruit (not juices)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Beef or pork hot dogs, regular-fat	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Cheese or cheese spread, regular-fat	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
French fries, home fries, or hash brown potatoes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Margarine or butter on vegetables, including potatoes	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Mayonnaise, regular-fat	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Salad dressings, regular-fat	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Rice	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Margarine, butter, or oil on rice or pasta	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

2. Over the past 12 months, when you prepared foods with margarine or ate margarine, how often did you use a reduced-fat margarine?

DIDN'T USE MARGARINE     
  Almost never     
  About 1/4 of the time     
  About 1/2 of the time     
  About 3/4 of the time     
  Almost always or always

3. Overall, when you think about the foods you ate over the past 12 months, would you say your diet was high, medium, or low in fat?

High     
  Medium     
  Low

**Reference:**

Thompson, F. E., Kipnis, V., Subar, A. F., Schatzkin, A., Potischman, N., Kahle, L., McNutt, S. (1998). Performance of a short instrument to estimate usual dietary intake of percent calories from fat. European Journal of Clinical Nutrition, 52(Suppl 2), S63 [Abstract].